(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 14 September 2006 (14.09.2006)

PCT

(10) International Publication Number WO 2006/094387 A1

(51) International Patent Classification:

A01N 25/16 (2006.01) C11D 17/00 (2006.01) A01N 55/10 (2006.01) A01N 31/02 (2006.01)

C11D 3/48 (2006.01)

(21) International Application Number:

PCT/CA2006/000320

(22) International Filing Date: 7 March 2006 (07.03.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/658,580

7 March 2005 (07.03.2005) US

(71) Applicant (for all designated States except US): ETHENA HEALTHCARE INC. [CA/CA]; 76 Adams Blvd., Brantford, Ontario N3S 7V2 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KOIVISTO, Bruce, Michael [CA/CA]; RR 1, Willsonville, Ontario N0E 1Z0 (CA). FERNANDEZ DE CASTRO, Maria, Teresa [CA/CA]; 67 Hobart Crescent, Brantford, Ontario N3P 1P7 (CA). MUNOZ, Francisco [MX/CA]; 67 Hobart Crescent, Brantford, Ontario N3P 1P7 (CA).

(74) Agent: HILL & SCHUMACHER; 87 Falcon Street, Toronto, Ontario M4S 2P4 (CA).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HIGH ALCOHOL CONTENT FOAMING COMPOSITIONS WITH SILICONE-BASED SURFACTANTS

(57) Abstract: This invention relates to a ôhigh lower alcohol contentő(>40% v/v of a C1-4 alcohol) liquid composition able to be dispensed as a stable foam with the use of non-propellant foam dispensing devices from non-pressurized containers. The liquid compositions comprise an alcohol, C1-4 (>40% v/v), a silicone-based surfactant of at least 0.001% by weight to prepare a foamable composition, 0-10% w/w of additional minor components added to obtain the desired performance (a foamable composition), and the balance being purified water. The compositions may include emulsifier-emollients and mosturizers, secondary surfactants, foam stabilizers, fragrances, antimicrobial agents, other type of medicinal ingredients, and the like ingredients or additives or combinations thereof commonly added to alcohol gels or foams, aerosol compositions or to toiletries, cosmetics, pharmaceuticals and the like.

2006/094387

HIGH ALCOHOL CONTENT FOAMING COMPOSITIONS WITH SILICONE-BASED SURFACTANTS

CROSS REFERENCE TO RELATED U.S. APPLICATIONS

5

This patent application relates to, and claims the priority benefit from, United States Provisional Patent Application Serial No. 60/656,580 filed on March 7, 2005, in English, entitled HIGH ALCOHOL CONTENT FOAMING COMPOSITIONS WITH SILICONE-BASED SURFACTANTS, and which is incorporated herein by reference in its entirety.

10

15

FIELD OF THE INVENTION

The present invention relates to compositions with high contents of lower alcohol (C_{1-4}) that could be dispensed as a foam both under low pressure from unpressurized containers and from pressurized containers achieved with an aerosol packaging system. The compositions to be dispensed as foams contain a silicone-based surfactant and when mixed with air provide a stable alcohol foam which can be used for personal cleaning or for disinfecting purposes.

.20

25

BACKGROUND OF THE INVENTION

Ethanol and/or Isopropyl alcohol and/or n-propyl alcohol compositions with at least 60% percent v/v (approximately 52% by weight) are well known to be antibacterial, therefore widely accepted for disinfecting purposes.

Nonetheless due to the inherent characteristics of alcohol, it is perceived that the higher the content the better the product and a solution with higher than 60% by volume alcohol content is more desirable.

Alcohol disinfectant solutions are generally thickened in order to eliminate the waste and facilitate spreading the composition throughout the desired area. It is also known that other than gelling agents one can use paraffin or waxes to achieve thickening of a solution with high alcohol concentration. Such a composition with landlin added to reduce the melting point closer to body temperature is described in United States Patent No. 2,054,989. One of the disadvantages of gels and the above-mentioned type of thick alcohol-containing compositions is that if they do not leave a tacky feeling on the hands after one use (although some do), the effect builds up after repetitive use during the day, making it necessary to eventually wash off the thickeners before continuing the usage of an alcohol antiseptic solution. The present invention if formulated for the above-mentioned type of product does not leave such a feel, and does not need to be washed off after repeated use.

15

20

25

10

.5 .

Generally speaking a high alcohol content disinfectant solution disinfects but does not clean. In order to make them disinfect and clean, so much soap would need to be added to the solution that the skin would feel soapy and disagreeable resulting in a formulation that would have little commercial appeal. Nonetheless, a non-irritant skin disinfecting formulation with a high content of a lower alcohol for use as a skin-washing agent has been successfully achieved by combining emulsifiers, surfactants and skin emollients as described in United States Patent No. 5,629,006.

Surfactants other than for cleaning purposes are also used for spreading an aqueous composition containing one or more active substances rapidly and evenly over a surface due to their wetting properties. The use of

good wetting agents definitely improves the efficient use of active substances in different compositions as described in United States Patent No. 5,928,993.

Although a high alcohol content disinfectant solution has good disinfectant characteristics, it has a sharp smell and is generally perceived to cause drying of the skin, characteristics which can also be diminished to a desirable level in the present invention.

A foam product with greater than 40% v/v alcohol, which is easy and safe to use, is desirable over conventional liquid, gel or ointment type composition products. The concentration of alcohol already poses a hazard in itself, and there are many applications in which the perceived risk may be diminished if it could be dispensed as a foam. A foam intended to be useful as a skin disinfecting agent must have a uniform consistency, spreadability, cleansing ability, and have a pleasant feel, i.e. have rapid breaking power when pressure is applied; all of which present a challenge for a high lower alcohol content composition.

The description of an aqueous foaming skin disinfecting composition using 15% w/w alcohol as a co-solvent, which requires no pressurized container or added propellant to produce the foam, is described in United States Patent No. 3,962,150.

20

5

10

15

The foam-forming agents utilized heretofore, have been incapable of forming stable foams when the liquid phase has high alcohol content without using other ingredients. Furthermore, lower alcohols have been considered to be defoamers rather than foam-promoting chemicals. According to Klausner, in United States Patent No. 3,131,153, if more than 64% alcohol is used non-

5

10

15

20

homogeneous compositions are obtained. The compositions in the patent required propellant to foam and the foams produced were of limited stability.

Various examples of compositions with a high content of a lower alcohol that are dispensed as a foam have been described, although for the purpose of the present invention the concentrations of alcohol and the levels of other ingredients are not suitable. More importantly, the use of propellants and aerosol containers to generate the foam is not desirable. For example, the compositions described in United States Patent No. 5,906,808 disclose a product that uses an emulsifying wax NF, and a combination of stearyl and cetyl alcohol, or other wax combinations, which improve the foaming performance of the composition, in combination with cetyl lactate, to produce a 0.8% chlorhexidine gluconate alcohol product.

United States Patent No. 5,167,950 issued to Lins discloses a foam product which requires a propellant and where no surfactant is added as a cleaning agent. The composition disclosed in this patent is based upon using an emulsifier system (fatty alcohol ROH 16-22 carbons) in combination with the use of a thickening agent (carbomer, klucel, etc.) to produce an antimicrobial aerosol mousse having a high alcohol content. The mousse includes alcohol, water, a polymeric gelling agent and a surfactant system comprising a C16-C22 alcohol, aerosol propellant and a non-ionic polyethoxylated surfactant.

Despite the work done to date it has been shown that there is little specific knowledge on how foams react and are formed, and surprisingly formulations that might seem not foamable result in the best foam producing ones while other formulations which seemed to have been producing foam

.

25

even while being prepared did not perform well at all in some non-aerosol foam dispensers. The behaviour of aqueous foams is not the same as that of an alcohol foam.

Silicone-based surfactants have been used in applications requiring lowering of the surface tension and increased wetting properties, especially in applications that require materials to be compatible with solvent systems other than water and non-reactive to other components in the compositions.

Silicone surfactants are desirable since they can achieve relatively low surface tension levels with relatively low concentrations in the compositions of interest Commercial examples of the exploitation of the advantage of the low surface tension levels achieved using Silicone based surfactants are crop protection products, printing inks, paints, floor coatings, etc. The characteristics mentioned above make silicone surfactants a candidate for the intention of this patent.

15

. 20

10

5

It would be very advantageous to have alcohol based disinfecting formulations containing silicone-based surfactants which may be dispensed as a foam under low pressure conditions and/and or through an aerosol packaging system. Further, it would be very advantageous and desirable to find a foaming agent that could be used in concentrations that would allow it to be used in products that can remain in the area on which they have been applied and do not need to be rinsed or wiped off due to small amounts of residue remaining after evaporation. Thus it would also be very advantageous to provide foams that do not leave an unpleasant sticky after-feel as most commercial alcohol gel products are known to, or which clog up the dispensing equipment used to dispense the foams. Silicone-based surfactants

25

are more than desirable for the purpose aforementioned since they are currently used as desirable cosmetic ingredients in creams, lotions, and other cosmetics due to their soft after-feel and properties.

5

10

15

20

25

SUMMARY OF THE INVENTION

The present invention provides high alcohol content compositions, which contain a surfactant/cleaning agent as well as a disinfectant/cleaning/solvent/carrier that causes very little drying to the skin or the hands of the user and is able to be dispensed as a foam from both pressurized and non-pressurized systems dispensing systems.

The present invention provides high alcohol content compositions that are able to be dispensed as a foam which are readily spread over the desired surface for the particular application. The present compositions can be formulated as an antimicrobial alcohol foam. The foamable compositions when dispensed from a suitable dispenser are stable and do not require the use of propellants and pressurized containers although if used would also foam.

Accordingly, the present invention provides foamable alcohol composition, comprising;

- a) a C₁₋₄ alcohol, or mixtures thereof, present in an amount greater than about 40% v/v of the total composition;
- b) an effective physiologically acceptable silicone-based surface active agent, which includes a lipophilic chain containing a silicone backbone ($-(R_2Si-O)_n)$, for wetting and foaming present in an amount of at least 0.01% weight percent of the total composition: and

c) water present in an amount to balance the total composition to 100% weight percent.

In this aspect of the invention the effective silicone-based surface active agent is present in an amount from about 0.001% to about 10.0% weight percent of the total composition which is physiologically acceptable so it can be used in personal care type products.

5

10

15

20

25

In a preferred embodiment of the invention the silicone-based surfactant may be a Bis-PEG-[10-20] dimethicone, a 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, an ethoxylated silicone-based surfactant, a Bis-PEG/PPG 18/6 Dimethicone, a Polyether-modified polysiloxane or a Polysiloxane betaine, or mixtures thereof. The present invention also provides an alcohol disinfecting composition, comprising:

- a) a) an alcohol C_{1-4} , or mixtures thereof, present in an amount between about 60% to about 80% v/v of the total composition;
- b) an effective physiologically acceptable silicone-based surface active agent for wetting and foaming present in an amount from about 0.01% to about 10.0% weight percent of the total composition;
- c) a foam stabilizing agent present in an amount from about 0.01 to about 12.0 % weight percent;
- d) any one of moisturizers, emollients and combinations thereof present in an amount from about 0.05 to about 5.0 % weight percent; and
- e) water in an amount to balance the total composition to 100% weight percent.

The present invention also provides a composition concentrate, comprising;

a) an effective silicone-based surface active agent for wetting and foaming present in an amount of at least 0.01% weight percent of the total composition 0.01to about 15.0 %;

- b) a foam stabilizing agent including at least from about 0.01 to about 10.0%;
- c) one of moisturizers, emollients and combinations thereof present in a range from about 0.05% to about 5.0%; and
 - d) water.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

5

10

15

20

25

The term "emollient" as used herein refers broadly to materials which are capable of maintaining or improving the moisture level, compliance, or appearance of the skin when used repeatedly.

The term "emulsifier" as used herein refers to surfactants or other materials added in small quantities to a mixture of two miscible liquids for the purpose of aiding in the formation and stabilization of an emulsion.

The phrase "emulsifying ingredients" as used herein is synonymous with emulsifier defined above.

The term "emulsion" as used herein refers to a colloidal dispersion of one liquid in another.

The term "surfactant" as used herein is the widely employed contraction for "surface active agents" which is the descriptive generic term for materials that preferentially adsorb at interfaces as a result of the presence of both, lipophylic and lipophobic structural units, the adsorption generally

resulting in the alteration of the surface or interfacial properties of the system.

The term "silicone-based surfactant" as used herein refers to a surface active agent in which the lipophylic chain contains a silicone chain, also described as any organosilicon oxide polymer in which the structural unit is usually comprising: $-R_2Si-O)_n$ — where R is a monovalent organic radical, and which enables the composition in which it is contained to clean, wet and foam.

The phrase "foam stabilizer" as used herein refers to an additive that increases the amount or persistence of foam produced by a surfactant system.

10

5

The term "disinfect" as used herein means to destroy or reduce harmful microorganisms.

The phrase "physiologically acceptable" as used herein means a material that does not usually produce irritation or toxicity when applied to the skin and is acceptable by users to apply to human skin

15

20

"Foam" as used herein means a liquid and a gas mixed to form a mass of small bubbles that has a structure that lasts for a variable length of time.

A bubble is a cell of gas surrounded by a film of liquid.

The term "aerosol" as used herein means a package and delivery system, and the product delivered, in which a pressurized gas is used to force the product out for dispensing. The gas may or may not be dispensed with the product.

An "aerosol foam" is a foam dispensed out of an aerosol package and delivery system as defined above.

The phrase "low pressure" in the context of producing a foam as used herein means a pressure of around an atmosphere or less such as when

5

10

15

20

dispensing the foam from an unpressurized conainer. Typically when foams are dispensed from aerosol containers the foam is considered to be being dispensed under high "pressure" conditions.

The present invention provides foamable alcohol compositions which include silicone-based surfactants with high contents of lower alcohol (C₁₋₄) able to be dispensed as a foam under low pressure conditions from unpressurized containers and through an aerosol packaging system. The present foamable compositions when mixed with air deliver a stable foam to provide an alcoholic liquid solution which can be used for personal cleaning or for disinfecting purposes and which breaks on pressure application such as when a user rubs their hands or when applied over a surface. All percentages provided herein are based on the total weight unless otherwise indicated.

The alcohol used in the present invention is a lower hydrocarbon chain alcohol such as a C₁₋₄ alcohol. The preferred alcohol is chosen from ethanol, 2-propanol, or n-propanol, most preferably ethanol, well accepted by Health Care personnel as an adequate disinfectant at the right percentages. It will be appreciated by those skilled in the art that if the alcohol used in the formulation is ethanol or a combination of ethanol with one or more of the other C₁₋₄ alcohols the ethanol preferably will be properly denatured to meet the local regulations of the targeted markets, but for the purposes of this patent it will be referred just as ethanol without it being specific as to whether it has been denatured. The compositions may use a single alcohol or as mentioned a blend of two or more alcohols may comprise the alcohol content of the composition

A significant and very surprising achievement of the present invention is that compositions suitable for disinfecting have been made containing greater than 40% v/v alcohol and a silicone-based surfactant able to be dispensed as a cosmetically appealing foam from both, low pressure conditions and through an aerosol packaging system.

The use of a silicone-based surfactant is the key ingredient as the primary foaming agent in the compositions designed to foam. Silicone surfactants have various interesting properties such as leaving little residue, being able to function in harsh chemical and thermal environments; they have an unparalleled wetting power, characteristics that are in general better than those of traditional surfactants, they show better surface-active properties in organic solvents, and that have made them widely used for applications in coatings, oilfield, material finishes, cleaning, paints, pesticides application, etc.

15

20

10

5

Traditional surfactants have carbon chains as the hydrophobic portion of the molecule. The hydrophile that is added to the carbon chain will determine the solubility and the general class of surface active agents as anionic, cationic, non-ionic or amphoteric. The silicone-based surfactants suitable for the compositions disclosed herein may include, but are not limited to, phosphate esters, sulphates, carboxylates, Imidazole quats, Amino Quats, Alkyl Quats, Amino propionates, ethoxylates, glycerol esters, amine oxides, acetylenic alcohol derivatives, phosphates, carbohydrate derivatives, sulfonates, betaines, Isethionates, esters, polyamides, and hydrocarbon surfactants that have a silicone chain $-(R_2-Si-O)_n$ - and are compatible with the other components being used for a particular formulation.

25

From the different silicone surfactants commercially available, samples from different manufacturers and different chemical surfactant groups were evaluated. Particularly, the Silicone Polyethers, also known as dimethicone copolyols showed the best performance. Amongst them, samples of multipendant and linear-diffunctional showed activity but the later was found to give superior foam properties. This is interesting as well as unexpected since in most water based applications if one of the multi-pendant or linear-diffunctional silicone surfactatns produces foam in water based mixtures, the other does not perform as well. However this is in agreement with the observations that foaming alcohol compositions behave quite differently from foaming water-based compositions.

5

10

15

20

25

While trying different surfactants, mixtures of two or more were evaluated to find out whether there was any synergy identifiable to optimize usage and foam performance. While some synergies were identified, it was also found that particularly difunctional silicone surfactants Bis-PEG [10-20] Dimethicones were the best when used alone. The notation Bis-PEG-[10-20] means all the Bis-PEG compounds having from 10 to 20 repeating oxyethylene groups. This applies to all other constituents as well. Specifically Bis-PEG 12 dimethicone and/or Bis-PEG-20 dimethicone and/or Bis-PEG-17 dimethicone are preferred along with 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate, a polyether-modified polysiloxane and a polysiloxane betaine, showed promising results but not as good as the dimethicone ones.

In preferred embodiments of the compositions, the effective siliconebased surface active agent may be a physiologically acceptable Bis-PEG-[10-

20] dimethicone, a 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate, a Polyether-modified polysiloxane or a Polysiloxane betaine, or mixtures thereof from about 0.01% to about 10.0% weight percent of the total composition.

5

10

It was surprisingly found that despite the characteristics of siliconebased surfactants, there is little or no information on their use to produce a foamable product with high alcohol content under either, low pressure conditions or through an aerosol packaging system.

Furthermore, in order to obtain a high alcohol content product able to produce a foam even if no pressurized containers or propellants are used, surface tension values as low as possible are required so that the pressure required to produce such foam by hand pumps and mechanical means would be sufficient.

During the development of the present invention, it was unexpectedly found that a relatively stable quick breaking foam could be obtained when using just ethanol and the silicone-based surfactant with as high as 80%v/w while using traditional surfactants at higher percentage yielded results that could not be even slightly similar and no foam at all could be obtained.

20

15

In order to achieve a commercially suitable formulation (one that lasts long enough for the purpose of use in disinfecting applications)., reducing the amount of silicone-based surfactant used while using the assistance of other ingredients such as secondary surfactants, emulsifiers, foam stabilizers, fragrances, and the like ingredients employed in cosmetics, aerosols, toiletries, personal care, etc. is one of the approaches that were followed. One of the commercial products obtained uses emulsifiers and polyethoxylated

2.5

5

10

15

20

25

fatty acid surfactants disclosed in United States Patent Nos. 5,167,950 and 6,090,395, both incorporated herein by reference, while other examples use a combination of different foam stabilizers to achieve a similar result.

Examples of secondary surfactants that may be used in the present compositions include other silicons surfactants, fluorinated surfactants, alkylglucosides, a poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ester of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, a betaine, a sulfobetaines, imidazoline derivatives, aminoacid derivatives, lecithins, phosphatides, some amine oxides and sulfoxides and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent.

A preferred betaine is cocamidopropyl betaine. A preferred alkylglucoside is cocoglucoside. A preferred fluorinated surfactant is DEA C[8-18] perfluoroalkylethyl phosphate; another preferred fluorinated surfactant is Ammonium C[6-16] perfluoroalkylethyl phosphate, Preferred polyethoxylated fatty alcohols are polyethoxylated stearyl alcohol (21 moles ethylene oxide) and polyethoxylated stearyl alcohol (2 moles ethylene oxide), and a combination of these two.

The compositions may include an antimicrobial agent. The following antimicrobials are offered as non-limiting examples of suitable antimicrobials

5

10

15

20

for use in the present invention and may include chlorhexidine salt, iodine, a complexed form of iodine, parachlorometaxylenol, triclosan, hexachlorophene, a phenol, behenyl alcohol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, other quaternary ammonium salts and mixtures thereof.

A preferred antimicrobial agent in the present compositions is chlorhexidine gluconate (CHG) present in an amount between about 0.10 % to about 4.0 % weight percent. Another preferred antimicrobial agent is didecyl dimethyl diamonium chloride in an amount between about 0.05% to 5% weight percent. Another preferred antimicrobial agent is Benzalkonium chloride in an amount between about 0.05% to 5% weight percent. Another preferred antimicrobial is Behenyl alcohol between 0.05-15% weight percent.

If the amount of ingredients employed is little enough not to leave a tacky feeling after the composition evaporates after single or multiple uses, and this is achieved while maintaining at least 60% v/v ethanol or n-propanol concentration or 70% v/v isopropanol, then the composition would be ideal for use as an alcohol hand sanitizer/disinfectant foamable composition.

The addition of water to the alcohol produces a more stable foam while allowing to reduce the amount-of-silicone-based surfactant required to foam the product. For example, using 0.5 to 1.0% silicone-based surfactant with a 50 to 60% v/v alcohol water solution produces a stable foam that does not readily collapse and that produces a stable puff that does not fall even when inverted and does not collapse until pressure is applied (such as when rubbed in hands or on over a surface) to provide an alcoholic liquid solution, while

levels of up to 5% are required if the percentage of alcohol used is greater than 65%w/w.

The use of a mild non-irritant surfactant widely used in the cosmetic industry such as cocamidopropyl betaine or a fluorinated surfactant such as DEA C[8-18] perfluoroalkylethyl phosphate or Ammonium C[6-16] perfluoroalkylethyl phosphate as a secondary surfactant is more suitable to prepare the foamable hydroalcoholic composition of the present invention depending on the silicone-based surfactant being used.

5

10

15

20

25

In order to stabilize the foam, foam stabilizers, as well as emulsifying ingredients have been tried with good results in allowing the product to be dispensed as a foam even when no propellant and/or pressurized container systems are used.

Examples of compatible foam stabilizers that can optionally be employed include lactic acid esters of monoglycerides, cationic emulsifiers, triquaternized stearic phospholipid complex, hydroxystearamide propyltriamine salts, lactic acid monoglycerides, food emulsifiers such as glyceryl monostearate, Behentrimonium chloride, Cetrimonium chloride, propylene glycol monostearate, glycols, sodium stearoyl lactylate, silicone wax, an encapsulated oil, Microcapsule Mineral Oil.

A preferred foam stabilizer used in the present foamable compositions is cetyl betaine. Another preferred foam stabilizer is glycerine. Another preferred foam stabilizer is Cetrimonium chloride and also Behentrimonium chloride.

Examples of moisturizers and/or emollients which may be used in the present formulations include lanolin, vinyl alcohol, polyvinyl pyrrolidone and

polyols selected from the group consisting of glycerol, propylene glycol, glyceryl oleate and sorbitol, cocoglucoside or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, ceteareth 20, and combinations thereof, present in an amount up to about 5%. The compositions may include a lipid layer enhancer such as a mixture of alkylglucoside and glyceryl Oleate, or PEG-200 Hydrogenated Glyceryl Palmiate, or Dihydroxypropyl PEG-5 Linoleammonium Chloride or PEG-7 Glyceryl Cocoate.

5

10

15

20

The compositions are formulated to be dispensed as a foam from an unpressurized dispenser having a dispenser pump for mixing the composition with air and dispensing foam therefrom. Alternatively the composition may be packaged in an aerosol container and dispensed under high pressure conditions.

The present invention is unique in that it can be dispensed as a foam from such manual low pressure dispensing systems as described, however, the composition can alternatively be dispensed under high pressure systems as well.

The composition may include an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition for pressurized discharge of the foam. The aerosol propellant may include propane, carbon dioxide, butane, dichloro difluoro methane, dichloro tetra fluoro ethane, octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane. When stored in a metal container with propellant, the formulation may include a corrosion inhibitor such as sorbic acid, benzoic acid, potassium sorbate and sodium

benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.

The inventors have very surprisingly discovered that it is possible to produce alcohol foams which include a combination of both the silicone surfactants disclosed herein and fluorosurfactants disclosed in copending U.S. Patent Application Serial No. 10/952,474, filed: September 29, 2004 and PCT Publication WO 2005/030917 A1 with the resulting foam exhibiting an interesting synergy with respect to the foam quality.

5

10

15

20

25

While both types of surfactants in the extremes of the concentrations of alcohol covered produce more and better foam when the alcohol concentration is closer to 40% and produce a lower quality foam that does not last in concentrations higher than 80% it was very interestingly found that the foam quality observed with fluorosurfactants is different from that obtained using silicone based surfactants.

With the pumps used for the examples, compositions using fluorinated surfactants are as described as an aereated "quick" breaking foam disappearing within seconds depending on the amount of alcohol and percentage of surfactant used, while with the silicone based surfactants the foam has a distinctive different appearance, it appears to be more dense and sometimes looks more uniform giving the appearance to be "whiter" not necessarily because there might seem to be more bubbles but because the bubbles last longer, they are more stable.

The difference is better observed when comparing side-by-side samples containing the same percentage of alcohol, and while it is evident that when the foam is first dispensed it shows more volume and a "better"

5

10

15

20

quality appearance if a fluorinated surfactant is used, this visual advantage is observed only during the first few seconds since the foam quite rapidly breaks down to a smaller volume to more resemble the foam obtained with the silicone-based surfactant. The most important difference appears to be how long the foam containing only the silicone-based surfactant lasts compared to the time the foam lasts with a foam containing only the fluorinated surfactant.

It should also be noted that one of the main differences in using one surfactant or the other (fluorosurfactants versus silicone surfactants) is the fact that fluorinated surfactants can reduce the surface tension to lower levels than those achieved using silicone-based surfactants alone, therefore compositions with only silicone based surfactants generally need a higher percentage of the silicone based surfactants to achieve a similar result.

A significant advantage to mixing both types of surfactants in the same composition is that the silicone based surfactant gives a longer lasting effect while the fluorinated surfactant has a better tolerance at higher levels of alcohol hence when mixed the combination provides an improvement in foam quality greater than obtaining using either one individually under some conditions.

Combining-the-surfactants allows the achievement of the desired foaming effect without exceeding solubility limits of either surfactant while at the same time taking advantage of the particular characteristics of the foam differences produced by each one.

The combination of silicone-based surfactants and fluorinated surfactant is of increasing advantage as the alcohol concentration increases, which is desirable to improve the microbiological efficacy.

The following non-limiting examples are set forth to show for the various preferred embodiments and are not in any way to limit the scope of the present invention.

5

10

15

20

EXAMPLES

Examples 1 through 12 were prepared to illustrate the ability to produce alcohol-based disinfecting formulations which can be dispensed as foams using different surfactants and a solution of water and 50% ethanol. Examples 13 through 16 illustrate increasing concentrations of a silicone-based surfactant to produce foam with 40% ethanol. Examples 17 through 32 show increasing concentrations of ethanol with different silicone-based surfactants to produce foam. Examples 33 through 36 illustrate increasing concentrations of a silicone-based surfactant to produce acceptable foam at 62% ethanol. Examples 37 through 52 illustrate the ability to produce foam using different surfactants and a solution of 70%v/v Isopropanol. All parts and percentages are expressed by weight unless otherwise indicated.

Comparatively, it was also found that for instance, Cocamidopropyl betaine (CAPB) alone even at 40% ethanol and at 3% CAPB, was unable to produce as good results as those with 60%v/v ethanol, and Silicone-based surfactants using much less percentage (less than 1.0%). Cocamidopropyl betaine does not give any acceptable foam above that percentage of alcohol and the lower than 60% v/v alcohol content makes it inadequate for a sanitizing solution. Also the solution left an unacceptable feeling on the skin after the alcohol evaporated (i.e. a soapy sticky feeling) indicating high levels of surfactant. Advantageously, the afterfeel of the present compositions was

25

5

10

15

20

not only not soapy but actually pleasant makes the present invention suitable for many different applications.

The following examples were intended to evaluate the foaming ability of different groups of Silicon surfactants with different combinations of ethanol, n-propyl alcohol and Isopropyl alcohol. They were also designed to asses the effect in the after feel as well as in the foam quality of some emollients and humectants, lipids, and other cosmetic type desired ingredients to be used with a leave-on hand sanitizer product. Some ingredients with antibacterial properties were also added to again evaluate their effect on foam quality and after feel.

Examples 53 to 293 have total alcohol contents from 61%w/w to 75%w/w. The best foam was achieved with only ethanol as the alcohol, with the next best being the combination Ethanol and n-propyl alcohol using up to 10% n-propyl alcohol. The worst foams were those obtained using only Isopropyl alcohol and/or n-propyl alcohol. Also interesting is the fact that the surfactant that works the best with just ethanol is not the same that works the best with the other 2 alcohols tried. More specifically it was found that the PEG-17-Dimethicone as well as the Bis-PEG/PPG 18/6 Dimethicone work better than the Bis-PEG 12 —Dimethicone with just n-propyl alcohol and/or Isopropyl alcohol. While combining different silicone surfactants if different alcohols are used seems to show a synergy to boost the foam when the addition of another alcohol disrupts the foaming ability of a particular silicone surfactant in the end the preferred silicone for the combination of alcohols emollients and other ingredients lies amongst the Bis-PEG [10-20]

Combinations of silicone surfactants with other surfactants were also tried; various experiments showed that although it was possible to find acceptable foam quality improvement when using other surfactants, the soapy after-feel was unacceptable for a leave-on product for many surfactants.

However the compositions combining silicone surfactants with fluorosufactants mentioned above showed an improvement that very advantageously allows for the reduction in the percentage of silicone surfactant to optimize percentages in the formulation regarding cost and other efficacy desired outcomes.

10

5

While some cosmetic ingredients like Cocoglucoside Glyceryl Oleate appear to deteriorate the foam quality when present in the composition, if the after-feel effect is beneficial increasing then the foam quality may be improved by increasing the percentage of surfactant present and/or adding a foam stabilizer to the composition to improve the foam quality.

15

It was also interesting to find that while a higher percentage of silicone surfactant present in the composition has been observed to increase the quality of the foam, the increased amount of silicone surfactant present has to be considerable before an improvement is observed. For example, the foam quality achieved with 1% was not significantly better with 1.5% but was considerably better with 2% and the foam quality using 3% or 4% was very similar while 5% was much better than just 3%.

20

25

The quality of the dispensed foam can be dependent upon the characteristics of the pump mechanism used for dispensing the foam from a bottle. For instance the quality of the foam was better with the smaller shot size of 0.75ml than the bigger shot size of 1.5ml from a pump for the same

size bottle. The shot size is the amount of liquid dispensed when the pump is activated.

From the above examples it can be concluded that they are clearly denoting how the art of foaming alcohol is different form that of foaming water based composition in more than one aspect and that unexpected outcomes could arise depending on the combination of ingredients desired for the final formulation in combination with the type of foaming device/mechanism to use.

Below are some more specific examples for compositions following a formulation to produce alcohol/silicone-surfactant hand/skin sanitizing foamable compositions; more than one being a foamable disinfecting composition with only alcohol being the only disinfectant ingredient, while other foamable disinfecting compositions use an added antimicrobial such as Chlorhexidine Digluconate or Didecyl Dimethyl Diammonium Chloride, Benzalkonium Chloride, Behenyl alcohol, etc.

15 **EXAMPLE 294**

Alcohol hand sanitizing foamable disinfecting composition

- 0.01- 5.0 % * silicone-based surfactant (primary surfactant)
- 0.01 1.0% cocoamidopropylbetaine (secondary surfactant)
- 0.05 1.0% cetyl betaine (foam stabilizing agent)
- 20 0.10 1.5% emulsifier fatty alcohol ROH 16-22 carbons or combination that works well in a final formulation containing
 - 60-70% v/v ethanol

5

10

25 :

- Q.S. water
 - Preferably a Bis-PEG-[10-20] dimethicone, a 3-(3-Hydroxypropyl) heptamethyltrisiloxane, ethoxylated, acetate, a Bis-PEG/PPG 18/6

5

10

15

Dimethicone, acetate, a Polyether-modified polysiloxane or a Polysiloxane betaine, or mixtures thereof.

EXAMPLE 295

Alcohol hand sanitizing foamable disinfecting composition Concentrate

- 0.1-5.0 % * a physiologically acceptable silicone-based surfactant*;(primary surfactant)
 - 0.001 12.0% 1,3 Butyleneglycol, 2-Butoxyethanol, or glycerin (foam stabilizing agents)
 - 0.05 5.0% cocoglucoside, glyceryl oleate (moisturizers, emollients and the like)
 - 60-70% v/v ethanol, n-propanol, isopropanol or a combination thereof Q.S. water
 - Preferably a Bis-PEG-[10-20] dimethicone, 3-(3-Hydroxypropyl)heptamethyltrisiloxane, ethoxylated, acetate, a Bis-PEG/PPG 18/6
 Dimethicone, acetate, a Polyether-modified polysiloxane or a
 Polysiloxane betaine, or mixtures thereof.

EXAMPLE 296

Alcohol hand sanitizing foamable disinfecting composition

- 0.01- 5.0 % * silicone-based surfactant (primary surfactant)
- 20 0.01 1.0% fluorinated surfactant and or other silicone-based surfactants or mixtures (secondary surfactant/s)
 - 0.05 1.0% Cetrimonium Chloride (foam stabilizing agent)
 - 0.05 1.0% Behentrimonium Chloride (foam stabilizing agent)
 - 0.10 1.5% Dihydroxypropyl PEG-5 Linoleammonium Chloride, Glyceryl
- Oleate, PEG-200 Hydrogenated Glyceryl Palmate, Behenyl PG-

Trimomium chloride, PEG-7 Glyceryl Cocoate or combination of emollients, lipids, humectants that works well in a final formulation containing

1-10% n-propyl alcohol

5 60-70% v/v ethanol

10

25

- Q.S. water
 - Preferably a Bis-PEG-[10-20] dimethicone, a 3-(3-Hydroxypropyl)heptamethyltrisiloxane, an ethoxylated, a Bis-PEG/PPG 18/6
 Dimethicone, acetate, a Polyether-modified polysiloxane or a
 Polysiloxane betaine, or mixtures thereof.

EXAMPLE 297

Alcohol hand sanitizing foamable disinfecting composition

0.01- 5.0 % * silicone-based surfactant (primary surfactant)

0.05 – 1.0% Cetrimonium Chloride (foam stabilizing agent)

15 0.05 – 1.0% Behentrimonium Chloride (foam stabilizing agent)

0.10 – 1.5% Dihydroxypropyl PEG-5 Linoleammonium Chloride, Glyceryl Oleate, PEG-200 Hydrogenated Glyceryl Palmate, Behenyl PG-Trimomium chloride, PEG-7 Glyceryl Cocoate or combination of emollients, lipids, humectants that works well in a final

20 formulation containing

1-10% n-propyl alcohol

60-70% v/v ethanol

Q.S. water

 Preferably a Bis-PEG-[10-20] dimethicone, a 3-(3-Hydroxypropyl)heptamethyltrisiloxane, an ethoxylated, a Bis-PEG/PPG 18/6 Dimethicone, acetate, a Polyether-modified polysiloxane or a Polysiloxane betaine, or mixtures thereof.

EXAMPLE 298

Alcohol hand sanitizing foamable disinfecting composition

- 5 0.01- 5.0 % * silicone-based surfactant (primary surfactant)
 - 0.01 1.0% fluorinated surfactant and or other silicone-based surfactants or mixtures (secondary surfactant/s)
 - 0.05 1.0% Cetrimonium Chloride (foam stabilizing agent)
 - 0.05 1.0% Behentrimonium Chloride (foam stabilizing agent)
- 10 0.10 1.5% Dihydroxypropyl PEG-5 Linoleammonium Chloride, Glyceryl Oleate, PEG-200 Hydrogenated Glyceryl Palmate, Behenyl PG-Trimomium chloride, PEG-7 Glyceryl Cocoate or combination of emollients, lipids, humectants that works well in a final formulation containing
- 15 60-70% v/v ethanol

20

- Q.S. water
- Preferably a Bis-PEG-[10-20] dimethicone, a 3-(3-Hydroxypropyl)heptamethyltrisiloxane, an ethoxylated, a Bis-PEG/PPG 18/6
 Dimethicone, acetate, a Polyether-modified polysiloxane or a
 Polysiloxane betaine, or mixtures thereof.

EXAMPLE 299

Alcohol hand sanitizing foamable disinfecting composition

- 0.01- 5.0 % * silicone-based surfactant (primary surfactant)
- 0.05 1.0% Cetrimonium Chloride (foam stabilizing agent)
- 25 0.05 1.0% Behentrimonium Chloride (foam stabilizing agent)

0.10 – 1.5% Dihydroxypropyl PEG-5 Linoleammonium Chloride, Glyceryl Oleate, PEG-200 Hydrogenated Glyceryl Palmate, Behenyl PG-Trimomium chloride, PEG-7 Glyceryl Cocoate or combination of emollients, lipids, humectants that works well in a final formulation containing

60-70% v/v ethanol

5

10

20

Q.S. water

Preferably a Bis-PEG-[10-20] dimethicone, a 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, an ethoxylated, a Bis-PEG/PPG 18/6
 Dimethicone, acetate, a Polyether-modified polysiloxane or a Polysiloxane betaine, or mixtures thereof.

EXAMPLE 300

Chlorhexidine gluconate (CHG) & alcohol hand sanitizing foamable disinfecting composition

15 Formulation 294, 295, 296, 297, 298 or 299 added with 0.50 – 4.0% Chlorhexidine Gluconate (CHG)

EXAMPLE 301

Didecyl Dimethyl Diammonium Chloride & alcohol hand sanitizing foamable disinfecting composition

Formulation 294, 295, 296, 297, 298 or 299 added with 0.01 – 5.0% Didecyl Dimethyl Diammonium Chloride

EXAMPLE 302

Benzalkonium Chloride & alcohol hand sanitizing foamable disinfecting composition

Formulation 294, 295, 296, 297, 298 or 299 added with 0.01 – 5.0% Benzalkonium Chloride

EXAMPLE 300

Behenyl Alcohol & alcohol hand sanitizing foamable disinfecting composition

Formulation 294, 295, 296, 297, 298 or 299 added with 0.01 – 5.0% Behenyl Alcohol

5

10

15

20

herein is straightforward since most of the ingredients are liquid. When wax type ingredients are to be used, they can be incorporated by warming up to 40-45°C preferably to the water portion while mixing and then allowing it to cool down or they could be added in "cold", at room temperature to the alcohol before any other ingredient and mixed until completely incorporated before adding the rest of the ingredients according to the composition.

The process to prepare the compositions of the present invention

Active ingredients may be pre-dissolved into the water first, a process that will be well known to anyone skilled in the art. If a specific formulation cannot be adjusted to give the desired quality of foam with the preferred percentages of the different constituents, the characteristics of the dispensing mechanism may be modified to improve foam quality, for example the type and characteristics of the foaming pump, such as changing the air/liquid ratio, screen sizes at the nozzle, to mention a few, can be adjusted in ways which will be apparent to those skilled in the art.

The compositions described herein provide improved alcohol based disinfecting products over commercially available compositions with high concentrations of alcohol, as well as the fact they are able to foam without the use of propellants or pressurized containers, although it will be appreciated that using propellants may in some cases improve the quality of the resulting foam.

5

10

15

20

25

Depending on the alcohol concentration and the application of the particular composition the foam produced can widely vary, being at the high end of a relatively fast breaking foam variety which is stable enough to be thoroughly spread onto the skin without undue waste or effort.

The present formulations may be first made as a concentrate with only some of the constituents which can be shipped and then constituted with the remaining constituents. For example, the concentrate can include a) the effective silicone-based surface active agent for wetting and foaming present in an amount of at least 0.01% weight percent of the total composition 0.01to about 15.0%, a foam stabilizing agent including at least from about 0.01 to about 10.0%, and any one of moisturizers, emollients and combinations thereof present in a range from about 0.05% to about 5.0%; and water.

The composition concentrate can then be constituted as an alcohol disinfecting composition by adding a) an alcohol C_{1-4} , or mixtures thereof, present in an amount between about 60 % v/v to about 80% v/v of the total composition; and b) water present in an amount to balance the total composition to 100% weight percent.

Due to the nature of the base composition with respect to the alcohol concentration and the quality of the ingredients, an advantageous application

for the present invention is as an alcohol skin/hand disinfectant composition for a foamable product, examples of which are described above.

Nevertheless, the present invention lends itself to the preparation of a wide variety of products for disinfecting applications, and it is to be understood that

other embodiments may be utilized and that changes may be made without

departing from the scope of the invention.

foam, antiperspirant foam, hair conditioner foams.

Consequently, it is intended that the claims be interpreted to cover such modifications and equivalents. To note a few, the following products may be produced using the alcohol/silicone-surfactants: medicated foams, sunscreen foams, hand cream foams, brush-less shaving cream foams, shower or bath oil foams, dry hair shampoo foams, make-up remover foams,

analgesic foam rubs, hair grooming foams and antiperspirants hair cleaning

As used herein, the terms "comprises", "comprising", "includes" and "including" are to be construed as being inclusive and open ended, and not exclusive. Specifically, when used in this specification including claims, the terms "comprises", "comprising", "includes" and "including" and variations thereof mean the specified features, steps or components are included. These

terms are not to be interpreted to exclude the presence of other features,

20 steps or components.

5

10

15

25

The foregoing description of the preferred embodiments of the invention has been presented to illustrate the principles of the invention and not to limit the invention to the particular embodiment illustrated. It is intended that the scope of the invention be defined by all of the embodiments encompassed within the following claims and their equivalents.

REFERENCES CITED

U.S. PATENT DOCUMENTS

2,054,9899/1936 Moore	‡2 55 7
3.962.1506/1976 Leonard et al	‡2 55 7
	55 7
4,440,6534/1984 James et al252/8.	7
5 167 950 12/1992 Lins	
4,956,170 09/1990 Lee514/772	.1
5,629,0065/1997 Minh et al424/405	
5,906,8085/1999 Osborne, et al 424/43	3
5,928,9937/1999 Ingegärd504/116	
5,951,993 09/1999 Scholz et al424/40	5
6,090,395 07/2000 Asmus et al424/40	1
6,610,315 08/2003 Scholz et al424/41	5
6,623,744 09/2003 Asmus et al424/40	1
6,562,360 05/2003 Scholz et al424/40	5

OTHER PUBLICATIONS

Myers, Drew; "Surfactant Science and Technology", second edition, Drew Myers, VCH Publishers, New York, 1992

1 -6
es
虿
Ε
a
×

Amount							
Ingredients	EX. 1	Ex. 2	Ex. 3	Ex. 4	Ex. 5	Ex. 6	
Ethanol Silions based surface to the	50.00	50.00	20.00	50.00	50.00	50.00	
Oncorre-Dased surfactant	0.50						
Cocal illidopropyl betaine (1)		8.00			.		
Alkylglucoside (Z)			8.00		1		
Alkylgiucoside (3)	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			8.00			
Oction Commission		1			8.00	***************************************	
Solutian Sesquioleate				***************************************	ļ.	8.00	
Purified Water	49.50	. 42.00	42.00	. 42.00	42.00	42.00	•
Total 0/							
iotal %	1.00.00	100.00	100.00	100.00	100.00 100.00 100.00	100.00	

(1) Amphoteric, (2) Nonionic, (3) Anionic

Examples 7-12

Amount Ingredients	Ex. 7 E	Ex. 8	Ex. 9	Ex. 10	Ex. 10 Ex. 11	Ex. 12
Ethanol Polysorbate 20 Polyoxyethylene Sorbitan Monooleate Sorbitan Monooleate Cocamidopropyl betaine & sodium caproyl lactate Cocamidopropyl hydroxysultaine Sodium Cocoamphoacetate Purified Water	50.00 8.00 8.00 42.00	50.00 8.00 42.00	50.00 8.00 42.00	8.00	50.00 8.00 42.00	50.00 8.00 42.00
Total %	100.00 16 Examp	100.00 Examples 13-16	100.00	100.00	100.00 100.00 100.00	100.00
Amount Ingredients		Ex. 13	Ex. 13 Ex. 14	Ex. 15 Ex. 16	Ex. 16	ŀ
Ethanol Bis-PEG-20 dimethicone 3-(3-Hydroxypropyl)-heptamethyltrisiloxane,ethoxylated, acetate Polyether-modified polysiloxane Polysiloxane betaine Purified Water	ethoxylated, acet	40.00 0.01 ate 59.99	0.01	0.01	40.00 0.01 59.99	ı
Total %		100.00	100.00 100.00	100.00	100.00	i

Examples 17-20

Ethanol Bis-PEG-20 dimethicone Durified Water	40.00 0.01 59.99 100.00	50.00 0.01 49.99 100.00	60.00 0.01 39.99 100.00	75.00 8.00 17.00	8.6
Total %	100.00	100.00	100.00	12/	3 8;
					100.00
	Examples 21-24	s 21-24			
Amount Ingredients	Ex. 21	Ex. 22	Ex. 23	ŭ	Ex. 24
Ethanol Polysiloxane betaine Purified Water	40.00 1.0 59.00	50.00 1.0 49.00	60.00	8 8 7.1	75.00 8.00 17.00
Total %	100.00	100.00	100.00	100	100.00

5-28
les 2
Examp

Amount Ingredients Ex.	Ex. 25	Ex. 26	Ex. 27	Ex. 28
Ethanol Polyether-modified polysiloxane 1 Purified Water 59	40.00 1.0 59.00	50.00 1.0 49.00	60.00 1.0 39.00	75.00 1.0 24.00
Total % 100	100.00	100.00	100.00	100.00
Example	Examples 29-32			
Amount Ingredients	Ex. 29	Ex. 29 Ex. 30	Ex. 31 Ex. 32	Ex. 32
Ethanol 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate 0.5 Purified Water	40.00 ate 0.5 59.5	50.00 0.5 49.50	60.00 0.5 39.50	75.00 10.00 15.00
Total %	100.00	100.00 100.00	100.00 100.00	100.00

9
3
ന
m.
S
a)
_
Q
_
_
Œ
×
ш

Amount Ingredients	Ex. 33	Ex. 34	Ex. 35	Ex. 36	· .
Ethanol	62.00	62.00	62.00	62.00	
Bis-PEG-20 dimethicone	0.50	1.00	2.0	5.00	
Purified Water	37.50	37.00	36.00	33.00	
Total %	100.00	100.00	100.00	100.00	

Examples 37-42

Amount Ingredients	Ex. 37	Ex. 38	Ex. 39	Ex. 40	Ex. 40 Ex. 41 Ex. 42	Ex. 42
70%v/v Isopropanol	99.90	92.00	92.00	92.00	92.00 92.00	92.00
Silicone-based surfactant	0.10					
Cocamidopropyl betaine (1)		8.00				
Alkylglucoside (2)	1	1	8.00	***************************************		1
) ap	-			8.00		1
Glycomul L		1			8.00	
Sorbitan Sesquioleate						8.00
		-				
Total %	100.00	100.00	100.00	100.00	100.00 100.00 100.00	100.00

(1) Amphoteric, (2) Nonionic, (3) Anionic

Examples 43-48

Amount Ingredients	Ex. 43	Ex. 44	Ex. 45	Ex. 46	Ex. 46 Ex. 47 Ex. 48	Ex. 48
70%v/v Isopropanol	92.00	92.00	92.00	92.00	92.00	92.00 92.00
Polysorbate 20	8.00				1	1
Polyoxyethylene Sorbitan Monooleate	1	8.00				
Sorbitan Monooleate			8.00		-	
Cocamidopropylbetaine & sodium caproyl lactate	1 lactate			8.00	1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Cocamidopropyl hydroxysultaine	1				8.00	***************************************
			1	1.	•	8.00
Total %	100.00	100.00	100.00	100.00	100.00	100.00 100.00 100.00

Examples 49-52

Amount Ingredients	Ex. 49	Ex. 50	Ex. 49 Ex. 50 Ex. 51 Ex. 52	Ex. 52
70%v/v Isopropanol Bis-PEG-20 dimethicone 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate Polyether-modified polysiloxane Polysiloxane betaine	1.0	99.00 99.00	99.00 99.00	99.00
Total %	00.00	100.00	100.00 100.00 100.00 100.00	100.00

The solutions prepared, were evaluated as to whether foam was produced or not and if so, then the foam produced was described as follows:

	SS		secs	secs	ninute		secs		nute	m	ninute		secs												secs	secs	secs	45secs
	Runny watery foam which lasts more than 20secs	Runny foam which lasts more than 20secs	Quick fast breaking foam lasts more than a 45secs	Quick fast breaking foam lasts more than a 45secs	Very good creamy and soft lasts more than a minute	Good creamy and soft lasts more than a minute	Quick fast breaking foam lasts more than a 45secs		Quick fast breaking foam lasts more than a minute	 Good creamy and soft lasts more than a minute 	Very good creamy and soft lasts more than a minute	Very good creamy and soft lasts minutes	Quick fast breaking foam lasts more than a 20secs												Runny watery foam which lasts more than a 45secs	Runny watery foam which lasts more than a 45secs	Runny watery foam which lasts more than a 45secs	Runny watery foam which lasts more than a 45
20	Yes	Yes	Yes	N _o	Yes	Yes	Yes	N	8	Š	No.	S S	Yes	8	Š	S N	8	S	8	<u>8</u>	Š	8	S N	2	Yes	Yes	Yes	Yes
EA. 24	Ex. 25	Ex. 26	Ex. 27	Ex. 28	Ex. 29	Ex. 30	Ex. 31	Ex. 32	Ex. 33	Ex. 34	Ex. 35	Ex. 36	Ex. 37	Ex. 38	Ex. 39	Ex. 40	Ex. 41	Ex. 42	Ex. 43	Ex. 44	Ex. 45	Ex. 46	Ex. 47	Ex. 48	Ex. 49	Ex. 50	Ex. 51	Ex. 52

Examples 53-63

INGREDIENTS	53	54	55	56	22	58	29	89	61	62	63
1,3-butylene glycol	0.43	0.43	-	-	1	0.43	0.43	-	-	-	0.43
Ethanol	62	25	52.2	62.2	42.2	62	52	52.2		42.2	62
Benzalkonium Chloride						0.1	0.1	0.1	0.1	0.1	
Dimethyl Ammonium											3
Chloride											- - -
PEG 17-Dimethicone	-	-	-	-	+	1	-	-	1	1	-
Glycerine	0.9	6.0	0.5	0.5	0.5	0.9	6.0	0.5	0.5	0.5	6.0
Cocoglucoside & Glyceryl Oleafe	0.08	90.0				0.08	0.08				0.08
DEA C8-C18						T					
Perfluoroalkylethyl	-					•					
Phosphate						-					
Ammonium C6-C16											
Perfluoroalkyfethyl	0.5	0.5	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5
Phosphate											
Cetrimonium Chloride											
n-Propyl alcohol	2	10	10		20	2	10	10	62.2	50	2
Propylene glycol			6.0	0.5	6.0			0.5	6.0	9'0	
Purified Water	33.1	36.1	34.6	34.6	34.6	33.2	35.2	34.5	34.5	34.5	33

Examples 64-73

INGREDIENTS	64	65	99	29	89	69	2	7.	22	73	64
1,3-butylene glycol	0.43	0.43	٦	-	_	0.43	0.43	0.43	0.43	0.43	0.43
	62	62	52.2	52.2	52.2	. 62	52	62	65	62	62
Benzalkonium Chloride											
Dimethyl Ammonium Chloride	0.5	1	0.1	0.1	-						0.5
PEG 17-Dimethicone	1	1	1	-	-	-	ر تن	-	-	-	1
	0.9	6.0	0.5	0.5	0.5	6.0	0.9	6.0	6.0	6.0	6.0
	0.08	80.0				0.08		0.08	0.08	90.0	0.08
							0.3	0.5	0.5	0.5	
_									!		
Ammonium C6-C16											
Perfluoroalkylethyl	0.5	0.5	0.3	0.3	0.3	0.5	0.3				0.5
				•							
Cetrimonium Chloride								0.05	0.1	0.05	
	2	2	10	10	10	2	10	2	2	2	2
			0.5	0.5	0.5						
	32.6	32.1	34.5	34.5	33.6	33.1	35	33	33	33	32.6

Example 74-84

														-			_		_
84	0.43	39	6:0	0.08	5.5							0.1					2		251
83	0.43	62	6.0	0.08	5.5							0.05					2		29
82	0.43	52	6.0	0.08		4.8			:		0.1						10		31.7
84	0.43	52	6.0	0.08		4.8					90'0						9		31.7
88	0.43	62	0.9	0.08	5.5						0.1						2		29
79	0.43	62	6.0	0.08	5.5			3	- - -		0.05						2 .		28.9
78	0.43	52	0.9	0.08		4.8		,									10		31.7
11	0.43	52	6.0	0.08		4.8		i C	 								10		31.7
92	0.43	62	6.0	0.08	5.5			č	 								2		29
75	0.43	62	0.9	90.0	5.5			<u> </u>	0.03								2		29
74	0.43	52	6.0	0.08		4.8	0.1				,						9		31.7
INGREDIENTS	1,3-butylene glycol.	Ethanol	Glycerine	Cocoglucoside & Glyceryl Oleate	Bis-PEG 17 Dimethicone	Bis-PEG 12 Dimethicone	Cetrimonium · Chloride	Dihydroxypropyl PEG-5	Linoleammonium	Chloride	Behenoyl PG- Trimonium Chloride	Behenamidopropyl Dimethylamine	Behentrimonium	BDLIQUIO	Cetearyl Alcohol and Repentrimonium	Chloride	n-Propyl alcohol	Propylene glycol	Purified Water

Example 85-94

INGREDIENTS	85	98	87	88	89	90	91	92	93	94
1,3-butylene glycol	0.43	0.43	0.43	0.43	0.43	0.43	0.43	0.43	0.43	0.43
Ethanol	52	52	62	62	52	52	62	62	25	52
Glycerine	6.0	0.9	0.9	0.9	0.9	0.9	0.9	6.0	6.0	0.9
Cocoglucoside & Glyceryl Oleate	. 0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08
Bis-PEG 17 Dimethicone			5.5	.5.5			5.5	5.5		
Bis-PEG 12 Dimethicone	4.8	4.8			4.8	4.8			4.8	4.8
Cetrimonium Chloride									·	
Dihydroxypropyl PEG-5										
Linoleammonlum Chloride								•		
Behenoyl PG- Trimonium Chloride										
Behenamidopropyl Dimethylamine	0.05	0.1								
Behentrimonium Chloride	:		0.5	0.1	0.05	0.1				
Cetearyl Alcohol and Behentrimonium Chloride			•		•		0.05	0.1	0.05	0.1
n-Propyl alcohol	10	10	2	2	10	10	2	2	10	5
Propylene glycol										
Purified Water	31.7	31.7	28.6	29	31.7	31.7	29	29	31.7	31.7

Examples 95-115

INGREDIENTS 95	95	96	97	86	66	100	191	102	103	104	105	198	107	108	109	110	111	112	113	114	115
Dimethicone PEG-8								1.5	2	-	1.5	2							-	1.5	2
Bis-PEG/PPG	,		Ç	,										,	,	Ι,	,	,	-		
Dimethicone	-	<u>.</u>	٧			٧							_	ပ်	·····	_	ე: 				
Ethanol													55	55	55	55	55	55	55	55	55
PEG-17										Ι,	,	١,									
Dimethicone										-	 - '										
Glycerine				-	-	T							r			-	-	-			Ī
Isopropyl Alcohol	20	70 70	70 70	70	20	02	22	20	22	02	02	22									
n-Propyl Alcohol									· ·			·	5	9	5	₽.	5	10	9	9	9.
Purified Water	29	28.5	28	28	27.5	27	29	28.5	28	88	27.5	27	34	33.5	33	33	32.5	32	34	33.5	33

Examples 116-136

	133 134 135 136	2 1 1.5 2
	132, 133	1.5
	129 130 131	
·		
_		
		1.5
_		-
_		1.5
		-
ç		18/6

Examples 137-156

			,							·			·		_	
156	<u> </u>					22				10	2	· ·				33
155						99				10	1.5					33.5
154.						22				10	1					34
153				က		61							Ò 4	- -		35.9
152	0.1			က		19										35.9
151		0.5		7.5		19										28
150				7		19										28
149				-		19										38
148				5	•					62			7	- - -		37.9
147	0.1			5						62						37.9
146				7.5						62			ć	- -		36.
145						22			ŀ	10	5	,				29
1						. 22			1	10	4					30
4 ₆	<u>.</u>					55			1	10	က					31
142	1	2							1	62						35
141		1.5							1	62						35.5
140		-							1	62						36
139		2	-							62				Ĺ		36
138		1.5								62						36.5
137		-								62						37
INGREDIENT S	Cetrimonium Chloride	Dimethicone PEG-8	Bis-PEG/PPG	18/6	 Dimethicone 	Ethanol	PEG-17	Dimethicone	Glycerine	n-Propyl Alcohol	Bis-PEG 12	Uimethicone	Dihydroxyprop yl PEG-5	Linoleammoniu	anionio iii	Purified Water

Examples 157-167

INGREDIENTS	157	158	159	160	161	162	163	164	165	166	167
1,3 Butylene Glycol											
Alcohol											
Behenamidopropyl Dimethylamine											
Behentrimonium Chloride							,				
Cetrimonium Chloride	0.1					0.1					-
Bis-PEG/PPG 18/6											
Dimethicone	_				_		•				
Ethanol	55	55	61	61	61	61	61				
Glycerine											
Cocoglucoside Glyceryl											
Oleate				-							
n-Propyl Alcohol	9	19						62	62	62	62
Bis-PEG 12 Dimethicone	1.5	1.5	-	1.5	7	1.5	1.5	-	1.5	7	1.5
Dihydroxypropyl PEG-5		.4									
Linoleammonlum Chloride		- - -					- -		-		
Purified Water	33.4	33.4	38	37.5	37	37.4	37.4	37 ·	36.5	36	36.4

Examples 168-177

NEDEDIENTS	168	169	170	171	172	173	174	175	1.76	177 ·
4 9 Butdone Glycol	2	43	43	4.3	4.3	4.3	4.3	4.3		
1,3 Butylene Clycol		62	63	62	62	69	62	.62		
Alcono		700	70	3	3	3				
Behenamidopropyl									0.1	
Dimethylamine										,
Behentrimonium Chloride	0.1									0.1
Cetrimonium Chloride										
Bis-PEG/PPG 18/6						•	7.5	7		
Dimethicone										į
Ethanol									5	٥
Chaorino		6.0	6.0	6.0	0.9	6.0	6.0	6.0		
CINCOLLIG										
Cocoglucoside Glyceryl		0.08	0.08	0.08	0.08	0.08	0.08	0.08		
Oleate						ľ	(
n-Propyl Alcohol	62	2	2	2	2	2	2	2	,	,
Bis-PEG 12 Dimethicone	1.5		-	1.5	2				7	7
Dihydroxypropyl PEG-5										
Linoleammonium Chloride								- 50	0.00	0
Purified Water	36.4	30.7	29.7	29.2	28.7	29.7	29.2	28.7	35.9	30.9

Examples 178-188

Examples 189-198

198				-				2				9		-						1	37
197								CA				55					13	•	-		33
196												61	2			-					37
195	0.43	2		-		0.1		2.5		`				6.0		0.08		0.5			25.5
194	0.43	20				0.1		2.5		;				6.0		0.08		0.5			25.5
193	0.43	70				0.1		•						6.0		0.08		0.5	2	2.3	25.5
192	0.43	20				0.1								6.0		0.08		0.1	2	7.3	25.9
191	0.43	62			0.1	0.1								6.0		0.08	2			7	32.4
190	0.43	62			0.1	0.1								6.0		90.0	2		r	2.5	31.9
189		55			0.1	0.1	2										10				32.8
INGREDIENTS	1,3-butylene glycol	Ethanol	Behenoyl PG-Trimonium Chloride	Cetearyi Alcohol	Behentrimonium Chloride	Cetrimonium Chloride	Dimethicone PEG-8	Bis-PEG/PPG 15/15	Dimethicone	Bis-PEG/PPG 18/6	Dimethicone	Ethanol	PEG-17 Dimethicone	Glycerine	Isopropyl Alcohol	Cocoglucoside & Glyceryl	n-Propyl Alcohol	PEG-200 Hydrogenated	Glyceryl Palmate	Bis-PEG 12 Ulmethicone	Purified Water

Examples 199-219

219						70	-							-	28
218						2								2	28
217	`					2							2		28
216						0.	2								82.
215						62	-		1	2		•	-		36
214						62				2				-	38
213						62	-			2				-	38
212						62				2		٠.		2	36
211						62				2			2		36
210						. 79	2			2					36
209						55	-			9			-		43
208						55				19			-	1	43
207						55	-			9				-	43
206						22				10				2	43
205						55				10			2		43
204						55	2			9		·			43
203	0.43	₩-	0.1			02		6.0	0.08					2.5	25
202				-	•	55	-			10					43
201	0.43		0.1			2		6.0	90.0		,			2.5	25
200						22	-	-		5				-	43
199	-					22	2			10				·	43
	1,3 Butylene Glycol	Cetearyl Alcohol and Behentrimonium	Chloride Behentrimonium Chloride	Bis-PEG/PPG 18/6	Dimethicone	· Ethanol	PEG-17 Dimethicone	Glycerine	Cocoglucoside Glyceryl Oleate	n-Propyl Alcohol	PEG-200 Hydrogenated	Glyceryl Palmate	Bis-PEG 20 Dimethicone	Bis-PEG 12 Dimethicone	Purified Water

Examples 220-230

INGREDIENTS 220 221 222 223	220	221	222	223	224	224 225	226		227 228	229	230
PEG-8			0.01	0 0	0.01 0.01 0.05	0.01	0.01	0.05	0.01	0.01	0.01
Dimethicone				5	20.5		5	3			
Ethanol	20	70	62	62	62	70	70	70	09	90	90
PEG-17											
Dimethicone		-									
n-Propyl Alcohol			2	2	2	2	2	2	10	10	10
Bis-PEG 20	,	*									
Dimethicone	-	-									
Bis-PEG-12	•		,	,	*	•	•	•	•	•	Ψ.
Dimethicone:	~ .		-	-	•	•	-	-	-	-	-
Purified Water	28	28	35	35	35	27	27	27	53	29	29

Examples 231-240

INGREDIENTS	231	232	231 232 233	234	234 235	236	236 237 238	238	239	240
PEG-8	200	2	900	2	2	200	5	5	200	5
Dimethicone	 	2.0	0.0	0.0	0.0	0.00	- O O	0.0		0.0
Ethanol				55	55	55	65	65	65	61
PEG-17										
Dimethicone										
n-Propyl Alcohol	9	09	09	10	10	10	10	10	10	39
Bis-PEG 20										
Olmethicone										
Bis-PEG-12	•	,	7	•	*		•		•	-
Dimethicone	-		-	-	-	-	•	-	-	-
Purified Water	39	39	39	34	34	34	24	24	24	7

Examples 241-251

		•	-	65			¥	-	-			10		23
7007		-	-	. 65			٠	-				10	1	22
7.49		•	-	09		•		-				10	1	27
248				62								2	-	32
247			•									2	1	27
246												09	-	39
245				61									;	38
244				92								10	7	24
243				55		•						10	τ-	34
242	90.0			61									1	38
241	0.01			61									-	38
INGREDIENTS	PEG-8- Dimethicone	Bis-PEG/PPG 18/6	Dimethicone	Ethanol	DEA C8-C18	Perfluoroalkylethyl Phosphate	Ammonium C6-C16	Perfluoroalkylethyl Phosphate	Cetrimonium Chloride	Dihydroxypropyl PEG-5	Linoleammonium Chloride	n-Propyl Alcahol	Bis PEG 12-Dimethicone .	Purified Water

Examples 252-261

		- i	_,						_			_
261			6				_	_				37
260			61				0.5	0.5				38
259			61				0.1	0.1				38.8
258			61				-				1	37
757			61				0.5	٠			1	37.5
526		•	61				0.1				1	37.9
255		7.5	65							10		23.5
254			65							10	1.5	23.5
253		-	65	-						10	1	22 ·
252		-	65	-						10		23
INGREDIENTS	PEG-8- Dimethicone	Bis-PEG/PPG 18/6 Dimethicone	Ethanol	DEA C8-C18 Perfluoroalkylethyl Phosphate	Ammonium C6-C16	Perfluoroalkylethyl Phosphate	Cetrimonium Chloride	Dihydroxypropyl PEG-5	Linoleammonium Chloride	n-Propyl Alcohol	Bis PEG 12-Dimethicone	Purified Water

Examples 262-272

INGREDIENTS	262	263	264	265	266	267	268	269	270	27.1	272
Behenyl PG-Trimonium Chloride											
Cetearyl Alcohol and Behentrimonium Chloride											
Polyethylene Glycol 600				0.1	0.5	-					
PEG-7 Glyceryl Cocoate.	-										
Ethanol	61	61	61	61	. 61	61	61	61	61	19	61
Cocoglucoside & Glyceryl										0.1	.0.5
Oleate											2
Polyethylene Glycol	0.2	0.5	,								
PEG-200 Hydrogenated							0.1	90	-		
Glyceryl Palmate								?:	•		
Bis-PEG 12 Dimethicone	1	_	, -	1	ļ	ı	ļ	.	1	1	7
Purified Water	37.8	37.5	37.	37.9	37.5	37	37.9	37.5	37	37.9	37.5

Examples 273-282

	-									
INGREDIENTS	273	274	275	276	277	278	279	280	281	282
Behenyl PG-Trimonium			•	0.1	6.0	+				
Cetearyl Alcohol and										
Behentrimonium Chloride							0.1	0.0		
Polyethylene Glycol 600										
PEG-7 Glyceryl Cocoate										0.1
Ethanol	61	61	61	61	.61	61	61	61	19	61
Cocoglucoside & Glyceryl										
Oleate	•									
Polyethylene Glycol				•						
PEG-200 Hydrogenated										
Glyceryl Palmate	•									
Bis-PEG 12 Dimethicone	-	2	က	γ-	1	1	·	1	.	-
Purified Water	37	37	36	37.9	37.5	37	37.9	37.5	37	37.9

Examples 283-293

INGREDIENTS	283	284	285	286	287	288	289	290	291	292	293
Behenyl PG- Trimonlum Chloride			0.2	0.25	0.2	0.25				0.2	0.2
Cetearyl Alcohol and Behentrimonlum			0.3	0.4	0.3	0.4				6.0	0.3
PEG-7 Glyceryl Cocoate	0.5	-	0.15.	0.2	0.15	0.2				0.15	0.15
PEG-PPG 20/6 Dimethicone									1		
PEG 4- Dimethicone							1				
PEG-8- Dimethicone			•					1	•		
Ethanol	61	61	61	61	65	65	65	65	92	61	91
Cocoglucoside & Glyceryl Oleate			0.2	0.25	0.2	0.25				0.2	0.2
Cetrimonium Chloride			0.1	0.2	0.1	0.2				0.1	0.1
Dihydroxypropyl PEG-5				i.		ų.				Č	. c
Linoleammonium Chloride			0.4	C:0	4					·	
n-Propyl Alcohol					10.	10	9	9	10		
PEG-200 Hydrogenated	٠		0.15	0.2	0.15	0.2				0.15	0.15
Glyceryl Palmate											
Bis-PEG 12 Dimethicone	-	1	2	. 2	7	ဇ				1.5	-
Purified Water	37.5	37	35.5	35	21.5	20	. 24	24	24	38	36.5

WHAT IS CLAIMED IS:

- 1. A foamable alcohol composition, comprising;
- a) a C₁₋₄ alcohol, or mixtures thereof, present in an amount greater than about 40% y/v of the total composition;
- b) an effective physiologically acceptable silicone-based surface active agent, which includes a lipophilic chain containing a silicone backbone $(-(R_2Si-O)_n -)$, for wetting and foaming present in an amount of at least 0.01% weight percent of the total composition; and
- c) water present in an amount to balance the total composition to 100% weight percent.
- 2. The composition according to claim 1 wherein the composition is foamable when mixed with air at low pressure, wherein when the composition is mixed with air, the mixture of the composition and air forms a foam.
- 3. The composition according to claims 1 or 2 wherein the effective silicone-based surface active agent for wetting and foaming is physiologically compatible, and wherein the silicone-based surfactant is selected from the group consisting of silicone ethoxylates, silicone glycerol esters, silicone amine oxides, silicone acetylenic alcohol derivatives, silicone carboxylates, silicone sulphates, silicone phosphates, silicone Imidazole quats, silicone Amino Quats, silicone phosphate esters, silicone carbohydrate derivatives, silicone Isethionates, silicone sulfonates, silicone betaines, silicone Alkyl

Quats, silicone Amino propionates, silicone esters, silicone polyamides, silicone fluorinated surfactants, and silicone hydrocarbon surfactants.

- 4. The composition according to claim 1, 2 or 3 wherein the effective silicone-based surface active agent is a physiologically acceptable silicone-based surface active agent selected from the group consisting of Bis-PEG-[10-20] dimethicones, 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate, polyether-modified polysiloxanes, polysiloxane betaine and mixtures thereof from about 0.01% to about 10.0% weight percent of the total composition.
- 5. The composition according to claim 1, 2, 3 or 4 wherein the alcohol C₁₋₄ is an aliphatic alcohol selected from the group consisting of methanol, ethanol, isopropanol, n-propanol, butanol and combinations thereof.
- 6. The composition according to claim 1, 2, 3, 4 or 5 wherein the silicone-based surface active agent is selected from the group consisting of Bis-PEG-12 dimethicone, Bis-PEG-17 Dimethicone, Bis-PEG-20 Dimethicone and combinations thereof.
- 7. The composition according to claim 1, 2, 3, 4 or 5 wherein the silicone-based surfactant is a 3-(3-Hydroxypropyl)-heptamethyltrisiloxane,
- 8. The composition according to claim 1, 2, 3, 4 or 5 wherein the siliconebased surfactant is a polyether-modified polysiloxane

9. The composition according to claim 1, 2, 3, 4 or 5 wherein the siliconebased surfactant is a polysiloxane betaine.

- 10. The composition according to claim 1, 2, 3, 4 or 5 wherein the silicone-based surfactant is a mixture of two or more of the silicone-based surfactants in claims 6, 7, 8 and 9.
- 11. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 wherein the alcohol is present in a range from about 40 % to about 90 % v/v.
- 12. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 wherein the alcohol is ethanol present in an amount of at least 60% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.
- 13. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 wherein the alcohol is a mixture of n-propanol and ethanol present in a combined amount of at least 60% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.
- 14. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 wherein the alcohol is a mixture of isopropanol and ethanol present in a combined amount of at least 60% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.
- 15. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 wherein the alcohol is isopropanol present in an amount of at least 70% v/v,

and wherein the composition is for use as an alcohol foam for personal hygiene.

- 16. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 wherein alcohol is n-propanol present in an amount of at least 60% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.
- 17. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 further including at least one additional surfactant for adjusting properties of the foam produced from the composition.
- 18. The composition according to claim 17 wherein the additional surfactant is selected from the group consisting of additional silicone-based surface active agents, fluorinated surfactants, alkylglucosides, poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ester of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, a betaine, a sulfobetaine, an imidazoline derivative, an aminoacid derivative, a lecithin, a phosphatide, an amine oxide, a sulfoxide and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent.

19. The composition according to claim 18 wherein the betaine is cocamidopropyl betaine.

- 20. The composition according to claim 19 wherein the alkylglucoside is cocoglucoside.
- 21. The composition according to claim 18 wherein the polyethoxylated fatty alcohol is polyethoxylated stearyl alcohol (21 moles ethylene oxide).
- 22. The composition according to claim 18 wherein the polyethoxylated fatty alcohol is polyethoxylated stearyl alcohol (2 moles ethylene oxide).
- 23. The composition according to claim 18 wherein the polyethoxylated fatty alcohol is a combination of polyethoxylated stearyl alcohol (21 moles ethylene oxide) and polyethoxylated stearyl alcohol (2 moles ethylene oxide).
- 24. The composition according to claim 18 wherein said fluorinated surfactant is DEA C[8-18] perfluoroalkylethyl phosphate.
- 25. The composition according to claim 18 wherein said fluorinated surfactant is ammonium C[6-16] perfluoroalkylethyl phosphate.

26. The composition according to claim 1, 2, 3, 4 or 5 wherein the silicone-based surface active agent is is 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate.

- 27. The composition according to any one of claims 1 to 26 including a foam stabilizing agent present in an amount up to about 10%.
- 28. The composition according to claim 27 wherein the foam stabilizing agent is selected from the group consisting of lactic acid esters of monoglycerides, cationic emulsifiers, quaternary ammonium compounds, triquaternized stearic phospholipid complex, hydroxystearamide propyltriamine salts, lactic acid monoglycerides, food emulsifiers such as glyceryl monostearate, propylene glycol monostearate, sodium stearoyl lactylate, cetyl betaine, glycolether, n-propanol, butyleneglycol, siliconee wax, an encapsulated oil, Microcapsule Mineral Oil, and combinations thereof.
- 29. The composition according to claim 27 wherein the foam stabilizing agent is selected from the group consisting of glycolether, glycerine, butyleneglycol, Behentrimonium chloride, or Cetrimonium Chloride and combinations thereof.
- 30. The composition according to any one of claims 1 to 29 including any one of a moisturizer, emollient, lipid layer enhancers and combinations thereof selected from the group consisting of lanolin, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol,

propylene glycol, butyleneglycol, glyceryl oleate and sorbitol, cocoglucoside, a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, ceteareth 20, an alkylglucoside, mixtures of alkylglucoside and glyceryl Cleate, PEG-200 Hydrogenated Glyceryl Palmiate, Dihydroxypropyl PEG-5 Linoleammonium Chloride or PEG-7 Glyceryl Cocoate, and combinations thereof, present in an amount up to about 5%.

- 31. The composition according to any one of claims 1 to 30 further comprising an acid or a base to adjust a pH of the composition to a preselected pH present in an amount from about 0.05 to about 0.5 % weight percent of the total composition.
- 32. The composition according to claim 31 wherein when an acid is used to adjust the pH the acid is selected from the group consisting of hydrochloric acid, citric acid and phosphoric acid, and wherein when a base is used to adjust the pH the base is sodium sesquicarbonate.
- 33. The composition according to any one of claims 1 to 32 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.
- 34. The composition according to any one of claims 1 to 33 including an antimicrobial agent.

35. The composition according to claim 34 wherein the antimicrobial agent is selected from the group of chlorhexidine salts, iodine, complexed forms of iodine, parachlorometaxylenol, triclosan, hexachlorophene, a phenol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, behonyl alcohol, and mixtures thereof.

- 36. The composition according to claim 34wherein the antimicrobial agent is chlorhexidine gluconate present in an amount between about 0.50 % to about 4.0 % weight percent.
- 37. The composition according to claim 34 wherein the antimicrobial agent is didecyl dymethyl diamonium chloride present in an amount between about 0:05 % to about 5.0 % weight percent.
- 38. The composition according to claim 34 wherein the antimicrobial agent is benzalkonium chloride present in an amount between about 0.05 % to about 5.0 % weight percent.
- 39. The composition according to claim 34 wherein the antimicrobial agent is behenyl alcohol present in an amount between about 0.05 % to about 5.0 % weight percent.
- 40. The composition according to any one of claims 1 to 39 further comprising constituents selected from the group consisting of organic gums

and colloids, lower alkanolamides of higher fatty acids, short chain diols and/or triols, fragrance, coloring matter, ultraviolet absorbers, solvents, suspending agents, buffers, conditioning agents, antioxidants, bactericides and medicinally active ingredients, and combinations thereof.

- 41. The composition according to any one of claims 1 to 40 stored in an unpressurized dispenser having a dispenser pump for mixing the composition with air and dispensing foam therefrom.
- 42. The composition according to any one of claim 1 to 40 stored in a pressurized dispenser having a dispenser pump for mixing the composition with air or a propellant and dispensing foam therefrom, the composition including an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition.
- 43. The composition according to claim 42 wherein the aerosol propellant is selected from the group consisting of propane, carbon dioxide butane, dichloro difluoro methane, dichloro tetra fluoro ethane octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane.
- 44. The composition according to claim 42 or 43 including a corrosion inhibitor is selected from the group consisting of sorbic acid, benzoic acid, potassium sorbate and sodium benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.

45. A composition concentrate, comprising;

a) an effective silicone-based surface active agent, which includes a lipophilic chain containing a silicone backbone ($-(R_2Si-O)_n -)$, for wetting and foaming present in an amount of at least 0.01% weight percent of the total composition 0.01to about 15.0 %;

- b) a foam stabilizing agent including at least from about 0.01 to about 10.0%;
- c) one of moisturizers, emollients and combinations thereof present in a range from about 0.05% to about 5.0%; and
 - d) water.
- 46. The composition concentrate according to claim 45 wherein the effective silicone-based surface active agent is a physiologically acceptable Bis-PEG-20 dimethicone, Bis-PEG-17 dimethicone, Bis-PEG 12 dimethicone, a 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate, a Polyether-modified polysiloxane or a Polysiloxane betaine, or mixtures thereof
- 47. The composition concentrate according to claim 45 or 46 wherein the foam stabilizing agent is selected from the group consisting of glycerine, cetrimonium chloride, behentrimonium chloride and combinations thereof.
- 48. The composition concentrate according to claim 45, 46 or 47 wherein the moisturizers and emollients include cocoglucoside, and glyceryl oleate, or PEG-200 Hydrogenated glyceryl palmiate, or Dihydroxypropyl PEG-5

linoleammonium Chloride or PEG-7 Glyceryl Cocoate, and combinations thereof, present in an amount up to 5%.

- 49. The composition concentrate according to claim 45, 46, 47 or 48 which is constituted as an alcohol disinfecting composition by adding
- a) an alcohol C_{1-4} , or mixtures thereof, present in an amount between about 60 % v/v to about 80% v/v of the total composition;
- b) water present in an amount to balance the total composition to 100% weight percent.
- 50. The composition concentrate according to claim 49 wherein the alcohol C₁₋₄ is selected from the group consisting of ethanol, n-propanol, isopropanol and combinations thereof.
- 51. The composition concentrate according to any one of claims 45 to 50 made by a process of mixing the constituents and then warming the concentrate to a temperature between about 30 to about 80 degrees Celsius prior to shipping the concentrate.
- An alcohol disinfecting composition, comprising;
- a) a) an alcohol C₁₋₄, or mixtures thereof, present in an amount between about 60% to about 80% v/v of the total composition;
- b) an effective physiologically acceptable silicone-based surface active agent, which includes a lipophilic chain containing a silicone backbone

 $(-(R_2Si-O)_n -)$, for wetting and foaming present in an amount from about 0.01% to about 10.0% weight percent of the total composition;

- c) a foam stabilizing agent present in an amount from about 0.01 to about 12.0 % weight percent;
- d) any one of moisturizers, emollients and combinations thereof present in an amount from about 0.05 to about 5.0 % weight percent; and
- e) water in an amount to balance the total composition to 100% weight percent.
- 53. The alcohol disinfecting composition according to claim 52 including air, wherein when the disinfecting composition is mixed with air, the mixture of disinfecting composition and air forms a foam.
- 54. The composition according to claims 52 or 53 wherein the effective silicone-based surface active agent for wetting and foaming is physiologically compatible, and wherein the silicone-based surfactant is selected from the group consisting of silicone ethoxylates, silicone glycerol esters, silicone amine oxides, silicone acetylenic alcohol derivatives, silicone carboxylates, silicone sulphates, silicone phosphates, silicone Imidazole quats, silicone Amino Quats, silicone phosphate esters, silicone carbohydrate derivatives, silicone Isethionates, silicone sulfonates, silicone betaines, silicone Alkyl Quats, silicone Amino propionates, silicone esters, silicone polyamides, silicone fluorinated surfactants, and silicone hydrocarbon surfactants.

55. The alcohol disinfecting composition according to claims 52 or 53 wherein the silicone-based surfactant is Bis-PEG-20 dimethicone.

- 56. The alcohol disinfecting composition according to claims 52 or 53 wherein the silicone-based surfactant is Bis-PEG-17 dimethicone.
- 57. The alcohol disinfecting composition according to claims 52 or 53 wherein the silicone-based surfactant is Bis-PEG-12 dimethicone.
- 58. The alcohol disinfecting composition according to claim 52 or 53 wherein the silicone-based surfactant is a 3-(3-Hydroxypropyl)-heptamethyltrisiloxane, ethoxylated, acetate,
- 59. The alcohol disinfecting composition according to claim 52 or 53 wherein the silicone-based surfactant is a Polyether-modified polysiloxane
- 60. The alcohol disinfecting composition according to claims 52 or 53 wherein the silicone-based surfactant is a Polysiloxane betaine.
- 61. The composition according to claims 52 or 53 wherein the siliconebased surfactant is a mixture of two or more of the silicone-based surfactants in claims 55 through 60.
- 62. The alcohol disinfecting composition according to any one of claims 52 to 61 wherein the foam stabilizing agent is selected from the group consisting

of lactic acid esters of monoglycerides, cationic emulsifiers, triquaternized stearic phospholipid complex, hydroxystearamide propyltriamine salts, lactic acid monoglycerides, food emulsifiers such as glyceryl monostearate, propylene glycol monostearate, sodium stearoyl lactylate, silicone wax, quaternary ammonium chlorides an encapsulated oil, Microcapsule Mineral Oil, butyleneglycol, and mixtures thereof.

- 63. The alcohol disinfecting composition according to any one of claims 52 to 62 wherein the foam stabilizing agent is selected from the group consisting of 2-butoxyethanol, glycerine butyleneglycol, Cetrimonium chloride, and or Behentrimonium chloride and combinations thereof.
- 64. The alcohol disinfecting composition according to claims 52 through 62 including any one of a moisturizer, emollient and combinations thereof selected from the group consisting of lanolin, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol, propylene glycol, butyleneglycol and sorbitol, or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, ceteareth 20, or an alkylglucoside and combinations thereof, present in an amount up to 5%.
- 65. The alcohol disinfecting composition according to claim 52 through 62 including a moisturizer and an emollient selected from the group consisting of glyceryl oleate, glycerine, cocoglucoside, or PEG-200 Hydrogenated Glyceryl Palmiate, or Dihydroxypropyl PEG-5 Linoleammonium Chloride or PEG-7

Glyceryl Cocoate.combinations thereof, present in an amount up to 5%.and combinations thereof.

- 66. The alcohol disinfecting composition according to any one of claims 52 to 62 including an antimicrobial agent.
- 67. The alcohol disinfecting composition according to claim 66 wherein the antimicrobial agent is selected from the group of a chlorhexidine salt, iodine, a complexed form of iodine, parachlorometaxylenol, triclosan, hexachlorophene, a phenol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, and mixtures thereof.
- 68. The alcohol disinfecting composition according to claim 66 wherein the antimicrobial agent is chlorhexidine gluconate present in an amount between about 0.50 % to about 4.0 % weight percent.
- 69. The alcohol disinfecting composition according to claim 66 wherein the antimicrobial agent is Didecyl Dymethyl Diamonium Chloride present in an amount between about 0.50 % to about 5.0 % weight percent.
- 70. The alcohol disinfecting composition according to claim 66 wherein the antimicrobial agent is Benzalkonium Chloride present in an amount between about 0.50 % to about 5.0 % weight percent.

71. The alcohol disinfecting composition according to claim 66 wherein the antimicrobial agent is Behenyl Alcohol present in an amount between about 0.50 % to about 5.0 % weight percent.

- 72. The alcohol disinfecting composition according to any one of claims 52 to 71 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.
- 73. The alcohol disinfecting composition according to any one of claims 52 to 71 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.
- 74. The alcohol disinfecting composition according to any one of claims 52 to 71 further comprising an acid or a base to adjust a pH of the disinfecting composition to a pre-selected pH present in an amount from about 0.05 to about 0.5 % weight percent of the total composition.
- 75. The alcohol disinfecting composition according to claim74 wherein the acid is selected from the group consisting of hydrochloric acid, citric acid and phosphoric acid, and the base is sodium sesquicarbonate.
- 76. The alcohol disinfecting composition according to any one of claims 52 to 71 stored in an unpressurized dispenser having a dispenser pump for mixing the disinfecting composition with air and dispensing foam therefrom.

77. The alcohol disinfecting composition according to any one of claims 52 to 71 stored in a pressurized dispenser having a dispenser pump for mixing the disinfecting composition with air or a propellant and dispensing foam therefrom, the disinfecting composition including an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition.

- 78. The alcohol disinfecting composition according to claim 52 wherein the aerosol propellant is selected from the group consisting of propane, carbon dioxide, butane, dichloro difluoro methane, dichloro tetra fluoro ethane octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane.
- 79. The alcohol disinfecting composition according to claim 77 or 78 including a corrosion inhibitor is selected from the group consisting of sorbic acid, benzoic acid, potassium sorbate and sodium benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.
- 80. The alcohol disinfecting composition according to any one of claims 52 to 79 including any one of a moisturizer, emollient and combinations thereof selected from the group consisting of lanolin, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol, propylene glycol, butyleneglycol and sorbitol, or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, ceteareth 20, or an alkylglucoside or PEG-200 Hydrogenated Glyceryl Palmiate, or Dihydroxypropyl PEG-5

Linoleammonium Chloride or PEG-7 Glyceryl Cocoate.combinations thereof, present in an amount up to 5%.and combinations thereof.

- 81. The alcohol disinfecting composition according to any one of claims 52 to 79 further comprising other constituents or materials including organic gums and colloids, lower alkanolamides of higher fatty acids, short chain diols and/or triols, fragrance, coloring matter, additional emollients, ultraviolet absorbers, solvents, emulsifiers, foam stabilizers or mixture of such stabilizers, suspending agents, buffers, conditioning agents, antioxidants, bactericides, medicinal active ingredient, and the like) that may be commonly added to aerosol compositions, toiletries, cosmetics and pharmaceuticals.
- 82. The composition according to any one of claims 1 to 44 further comprising a physiologically acceptable fluorosurfactant present in a range from about 0.01 to about 10%w/w, wherein said C_{1-4} alcohol is present in an amount from about 40 to about 90 v/v of the total composition, and wherein the silicone-based surface active agent for wetting and foaming present in an amount from about 0.1 to about 10%w/w weight percent of the total composition.
- 83. The composition according to claim 82 wherein the fluorosurfactant is selected from the group consisting of fluorinated ethoxylates, glycerol esters, amine oxides, acetylenic alcohol derivatives, carboxylates, phosphates, carbohydrate derivatives, sulfonates, betaines, esters, polyamides, silicones, and hydrocarbon surfactants.

84. The composition according to claim 82 or 83 wherein the fluorosurfactant is an amphoteric polytetrafluoroethylene acetoxypropyl betaine $(CF_3CF_2(CF_2CF_2)_nCH_2CH_2(OAc)CH_2N^+(CH_3)_2CH_2COO^-)$ where n=2 to 4.

- 85. The composition according to claim 82 or 83 wherein the fluorosurfactant is an ethoxylated nonionic fluorosurfactant of the following structure: $RfCH_2CH_2O(CH_2CH_2O)_xH$ where $Rf = F(CF_2 CF_2)y$, x = 0 to about 15, and y = 1 to about 7.
- 86. The composition according to claims 82 or 83 wherein the fluorosurfactant is an anionic phosphate fluorosurfactant of the following structure: $(RfCH_2CH_2O)_x P(O)(ONH_4)_y$ where $Rf = F(CF_2 CF_2)z$, x=1 or 2, y=2 or 1, x+y=3, and z=1 to about 7 or mixtures thereof.
- 87. A method of forming a foam using a composition according to any one of claims 1 to 44, by storing said composition in a container and having a dispenser pump activating the dispenser pump to mix the alcohol and silicone-based surface active agent mixture with air under low pressure to form a low pressure foam containing alcohol which is dispensed from said dispenser.

INTERNATIONAL SEARCH REPORT

International application No. PCT/CA2006/000320

A. CLASSIFICATION OF SUBJECT MATTER

 $\text{IPC: } \textit{A01N 25/16} \ (2006.01) \ , \ \textit{C11D 17/00} \ (2006.01) \ , \ \textit{C11D 3/48} \ (2006.01) \ , \ \textit{A01N 55/10} \ (2006.01) \ , \\$

A01N 31/02 (2006.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: A01N 25/16 (2006.01), C11D 17/00 (2006.01), C11D 3/48 (2006.01), A01N 55/10 (2006.01), A01N 31/02 (2006.01)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used)

Canadian Patent Database, Delphion full text search - silicone, foam, alcohol, surfactant, and - bis-PEG-20, 17, 12 dimethicone, USPTO - silicone, foam, alcohol, surfactant

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Х, Р	WO 05030917 (ETHENA HEALTHCARE INC. et al.) 7 April 2005 (07-04-2005) Claims 4-94.	1-3, 5, 11-23, 27-45, 47-54, 62-81, 87
x	US 5629006 (BECTON, DICKINSON AND COMPANY) 13 May 1997 (13-05-1997) Column 1 and 3, Claim 1.	1-5, 8, 11, 12, 14, 15, 17, 18, 27, 28, 30- 34, 40, 45, 46, 49, 50, 52-54, 59, 62-64, 66, 72-75, 80, 81
x	CA 2534692 (LRC PRODUCTS LIMITED) 17 February 2005 (17-02-2005) Page 2, lines 15-18, claims 1-28.	1-3, 5, 11, 12, 14, 17, 18, 28, 30-34, 40
		,

[]]	Further documents are listed in the continuation of Box C.	[X] See patent family annex.
* "A" "E" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier application or patent but published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention. "X" document of particular relevance; the claimed invention carnot be considered novel or cannot be considered to involve an inventive step when the document is taken almost document of particular relevance; the claimed invention carnot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
	of the actual completion of the international search une 2006 (13-06-2006)	Date of mailing of the international search report 30 June 2006 (30-06-2006)
Cana Place 50 V	e and mailing address of the ISA/CA adian Intellectual Property Office e du Portage I, C114 - 1st Floor, Box PCT fictoria Street neau, Quebec K1A 0C9	Authorized officer Sandra Nevill (819) 934-6732

Facsimile No.: 001(819)953-2476

والصحاف للأحاص

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No. PCT/CA2006/000320

Publication Date	Patent Family Member(s)	Publication Date	
07-04-2005	None		
13-05-1997	BR9502916 A	27-02-1996	
•			
•			
	EP0689767 A2	03-01-1996	
	JP2638575B2 B2	06-08-1997	
17-02-2005	AU2004262971 A1	17-02-2005	
	EP1663131 A1	07-06-2006	
	GB0416924D D0 WO2005013930 A1	01-09-2004 17-02-2005	
	Date 07-04-2005 13-05-1997	Date Member(s) 07-04-2005 None 13-05-1997 BR9502916 A	Date Member(s) Date 07-04-2005 None 13-05-1997 BR9502916 A 27-02-1996 CA2151774 A1 28-12-1995 DE69512841D D1 25-11-1999 DE69512841T T2 18-05-2000 EP0689767 A2 03-01-1996 JP2638575B2 B2 06-08-1997 17-02-2005 AU2004262971 A1 17-02-2005 EP1663131 A1 07-06-2006 GB0318160D D0 03-09-2003 GB0416924D D0 01-09-2004

THIS PAGE BLANK (USPTO)

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

efects in the images include but are not limited to the items checked:
BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)